Complex PTSD and Specialist Trauma Populations

THE TALKING THERAPIES UK PRACTITIONER SERIES



Complex PTSD and Specialist Trauma Populations

Developmental trauma, complex presentations and special contexts

Advanced level · 14 hours of continuing professional development

TALKING THERAPIES UK

First edition · 2026

Published by Talking Therapies UK, talkingtherapies.co.uk.

Copyright © 2026 Talking Therapies UK. All rights reserved.

The moral right of the author has been asserted. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior written permission of the publisher, except for brief quotations in a review, and except that a purchaser of the accompanying course may print a single copy for their own study.

First edition, 2026.

This manual accompanies the Complex PTSD and Specialist Trauma Populations course in the Talking Therapies UK Digital Classroom, from which its content is drawn and against which it is kept current.

Revision 5. This printing was set on 10 November 2026 from the course as it stood that day.

A note on the clinical material. Every case in this book is a fictional composite written for teaching. No real person appears in it, however disguised. Nothing here constitutes clinical supervision, and nothing in it replaces the judgement of a qualified practitioner working within their competence, their professional body's code and their local governance. Where guidance is cited it was current at the date of this edition; readers must check the standing of any guidance before relying on it.

Set in Poppins and Roboto, the Talking Therapies UK typefaces. Figures drawn by the Talking Therapies UK teaching-diagram engine.

Contents

How to use this manual **1**

Part 1 Foundations and Neurobiology of Complex PTSD **5**

1. Understanding complex PTSD 6
2. The phased approach 11

How to use this manual

This advanced training module provides an in-depth exploration of Complex PTSD and the nuanced presentations often arising from developmental trauma. Moving beyond standard PTSD, the curriculum delves into the DSM-5-TR and ICD-11 criteria, specifically focusing on the profound disturbances in self-organisation that characterise CPTSD. Learners will examine how early relational trauma disrupts the development of affect regulation, self-concept, and interpersonal functioning, gaining a comprehensive understanding of the neurobiological and psychological mechanisms underpinning these complex presentations.

A central component of this course is the detailed examination of the phased approach to treatment, a critical framework for managing the heightened risks associated with complex trauma. Participants will study the stabilisation phase, which establishes safety and containment, and the subsequent trauma processing stages, ensuring they are equipped to handle the intense emotional and physiological arousal that can arise during therapy. This section emphasises the importance of pacing and the use of evidence-based interventions tailored to the unique needs of clients with complex histories.

The course further extends its scope to specialist populations, offering specialised insights into the trauma experiences of military personnel, medical professionals, and first responders. By analysing the specific stressors inherent in these high-stakes environments, including moral injury and exposure to critical incidents, clinicians will learn to adapt their therapeutic approach to resonate with these specific cultural and occupational contexts. This module addresses the unique challenges of treating individuals whose professional roles involve high levels of responsibility and exposure to human suffering.

Additionally, the training addresses the intersection of trauma with relational dynamics and identity. Clinicians will explore how complex trauma impacts attachment styles, parenting capabilities, and a client's sense of self, providing the tools necessary to support clients in rebuilding healthy

relationships and integrating a coherent identity. The curriculum also covers the critical clinical skills required to manage dissociation and other complex symptom clusters, alongside essential guidance on maintaining therapist sustainability and preventing burnout when working with highly distressed clients.

This manual is the written form of a course taught in the Talking Therapies UK Digital Classroom. It is set at advanced level and carries 14 hours of continuing professional development for practitioners who complete the accompanying course and its assessment.

It is written to be read in order. Each part opens with what it is for; each chapter teaches one idea to a working depth and closes with the points worth keeping and a practice task that converts reading into something you do. The tasks are the part most often skipped and the part that most reliably changes practice, and several of them are designed to be taken to supervision rather than completed alone.

The figures are teaching diagrams rather than decoration. Each is the conventional form of a model you will meet in the literature and in supervision, drawn in one visual language across the whole series so that the shape of an idea is recognisable wherever it reappears.

Where a chapter draws on published work, the reference is named in the text and gathered in the further reading at the back. Terms of art are defined in the glossary, which is worth reading straight through once before you begin.

What this manual will leave you able to do

- Adapt trauma-focused work for complex PTSD: phasing, titration and relational repair
- Treat multiple-event and developmental trauma without endless stabilisation
- Work with dissociation, including flashback management and grounding protocols, at specialist level
- Address perpetrator-related presentations, moral injury and traumatic bereavement
- Adapt practice for refugees, veterans, emergency workers and abuse survivors

- Sustain yourself in high-exposure work: vicarious trauma as an occupational hazard to manage

PART 1

Foundations and Neurobiology of Complex PTSD

This module explores the core theory, neurobiological foundations, structural dissociation, and self-organisation disturbances inherent in complex PTSD.

CHAPTER 1

Understanding complex PTSD

Some trauma is an event; other trauma is an environment, and the difference is diagnostic. When the threat is prolonged or repeated, when escape is impossible or costs more than staying, captivity, trafficking, domestic violence, and above all abuse across a childhood, the injury extends beyond the fear machinery into the person's whole self-organisation, and ICD-11 gave that extension its own name. This lesson teaches complex PTSD as the distinct diagnosis it now is: what it adds to PTSD, why prolonged trauma produces the addition, and what the difference demands of treatment.

The architecture is core plus three. Complex PTSD requires the full PTSD core, re-experiencing in the here and now, avoidance, persistent current threat, and adds the disturbances of self-organisation, three domains in which the self, not just the alarm, was reshaped. Affect dysregulation: emotional responses that are too big or absent altogether, hair-trigger reactivity beside baffling numbness, feelings arriving as storms without names, calming taking hours where others take minutes, and, frequently, the desperate regulators that follow, self-harm, substances, dissociation. Negative self-concept: not the event-linked appraisals of simple PTSD, I was weak that night, but a settled identity, worthless, defiled, deserving, held with the certainty of something learned before words, and saturated with shame, which will be the treatment's constant companion. Relational disturbance: difficulty sustaining closeness of any kind, oscillation between hunger for connection and terror of it, trust extended too totally or refused entirely, and the repeating discovery that the strategies that survived the abusive relationship, appeasement, vigilance, pre-emptive abandonment, are dismantling every subsequent one.

Why does prolonged trauma produce this wider injury? Because chronicity converts responses into architecture. A single event is an exception the self survives; an environment is a curriculum the self is built inside. The threat system, activated not for minutes but for years, calibrates permanently high, and the developing stress axis calibrates with it, this is the developmental biology the personality disorder pathway teaches in depth, and the overlap is not accidental. Emotion regulation is not lost but never taught: the caregiver

who should have been the external regulator, soothing storms until soothing is internalised, was the storm's source, so the capacity has no ordinary history to restore, only a first history to build. Self-concept forms from the environment's mirror: a child treated as worthless does not conclude the environment is defective, children cannot afford that conclusion, they conclude they themselves are, Fairbairn's moral defence, it is safer to be a bad child in a world of good parents than the reverse, and the badness becomes load-bearing identity. Relational templates encode the only relationships available: love arriving with harm, care conditional on silence, attachment itself made dangerous, fright without solution carried forward as the adult's working model. And memory in such histories is often not one filed event but years of fragments, some pre-verbal, some dissociated at encoding, which is why the narrative work of the practitioner course will need the adaptations the next lesson teaches.

Differential discipline completes the picture. Complex PTSD versus PTSD turns on the self-organisation triad, present persistently and pervasively, not only in trauma-cued moments. Complex PTSD versus borderline-pattern personality disorder is the fine judgement the two pathways share: the presentations overlap substantially, dysregulation, unstable relationships, negative self-experience, and the ICD-11 discriminators are the required trauma core in complex PTSD, re-experiencing and avoidance must be present, and the characteristic colourings, fear of abandonment and identity instability oscillating around emptiness pointing towards the borderline pattern, stable worthlessness with trauma re-experiencing pointing towards complex PTSD, with both diagnoses sometimes true and co-recorded. The stakes of the judgement are practical: it routes people between this pathway's phased trauma treatment and the staged personality disorder frameworks, and mis-routing costs years. Depression, dissociative disorders and substance dependence co-travel routinely and are assessed, not assumed.

Set two histories side by side and the architecture teaches itself. The paramedic from the foundations course carries PTSD: a single catastrophic scene, intrusions with nowness, avoidance of one postcode, a hair-trigger startle, and beneath it a self that remains intact, she knows she is good at her job, loves clumsily but securely, and regulates well enough between symptoms; treat the memory and the meanings and her self-organisation will hold the recovery. Beside her, a woman of the same age referred with the

same word, trauma, after leaving a fifteen-year relationship that began as control and became violence, whose childhood had already taught the curriculum, a father whose rages set the house's weather, a mother who managed him by sacrificing the children's claims. She has the core: intrusions of specific assaults, avoidance of the town he lives in, the permanent scanning. And she has the triad: rage and numbness alternating without her consent; a certainty of her own worthlessness so old she calls it just knowing myself; and relationships, including with services, run on appeasement, vigilance and pre-emptive flight. Same referral word, different injuries: one is a wound in an intact person, the other is an architecture built under siege. The treatment consequences follow the architecture: the paramedic can move to memory work within weeks; the second woman needs the phased chassis, regulation built before processing, shame met as identity, the relationship itself doing years of the lifting. Diagnosis here is not labelling; it is load-bearing routing, and the six-line profile you are about to write is the routing instrument.

What the difference demands of treatment is the course's organising preview: the practitioner toolkit remains the engine, but it now needs a chassis built for load, longer work, phased structure, regulation capacities built before memory processing because they were never there to restore, shame addressed as identity rather than appraisal, and the therapeutic relationship carrying more of the treatment, because for injuries acquired in relationship, relationship is the only tissue that regrows. Close by applying the architecture: for a person you know, or a composite, write the six-line profile, two lines for the PTSD core as it shows in them, one for whether escape was possible, and one each for affect, self-concept and relationships, ending with the diagnostic sentence and one routing implication. The next lesson turns that profile into a treatment shape: the phased approach.

Going deeper: the diagnosis that reorganises the treatment.

ICD-11's complex PTSD is a structural claim, not a severity remark: alongside the PTSD core (re-experiencing, avoidance, current threat) stand three disturbances of self-organisation — affect dysregulation, negative self-concept, relational disturbance — that arise characteristically from prolonged, repeated or inescapable trauma, above all trauma inflicted early and relationally. The distinction matters because the DSO features are not decorations on PTSD; they are what happens when trauma occurs while the

regulatory and relational systems are still being built, or is administered by the very attachment figures those systems needed. The result is not only a set of unfiled memories but a development conducted inside threat: emotion regulation never scaffolded, a self-concept formed around the perpetrator's account, and relational templates in which closeness and danger share an address.

Clinically this reorganises everything. Differential diagnosis becomes consequential — CPTSD's overlap with borderline presentations is real, but the centre of gravity differs (a trauma-anchored, stable-negative self and avoidant relating, against identity instability and frantic attachment), and the treatment implications differ with it. Assessment must map dissociation carefully, because it is the population's signature regulation strategy. And treatment planning becomes phased by necessity rather than convention: the DSO features are both obstacles to memory work (a patient who dissociates at activation cannot process) and targets in their own right, which is why complex presentations receive stabilisation with exit criteria, titrated processing, and a therapy relationship treated as an instrument rather than a backdrop.

Hold the hopeful datum alongside the gravity: CPTSD responds to treatment. Longer, phased, relationally attentive treatment — but response, not management, is the evidence-based expectation.

A working note: the differential that matters most.

The complex-PTSD diagnosis shares its border with borderline-pattern personality difficulty, and the differential deserves a working method rather than an argument. Ask three questions of the presentation. Is there re-experiencing anchored to identifiable traumatic material? CPTSD requires it; personality-level difficulty does not. What is the shape of the relational disturbance: consistent avoidance and mistrust with a stable, negative self-account (CPTSD's signature), or oscillation, idealisation-devaluation, frantic responses to abandonment and an unstable self (the borderline signature)? And what happens under attachment stress: withdrawal and shutdown, or escalation towards the other person? Comorbidity is real and common, so 'both' is a legitimate answer, but the distinction still steers sequencing: CPTSD's disturbances are treated largely through and after memory work, while borderline-pattern difficulty usually needs its own stabilisation and relational treatment in which trauma processing is one

planned phase. Getting this differential right, in either direction, is worth more to the person than any subsequent technical excellence.




FIGURE 1.1 The window of tolerance

Siegel's window of tolerance describes the zone of arousal within which experience can be processed and integrated. Trauma narrows the window: survivors are thrown into hyperarousal (fight/flight) or hypoarousal (shutdown) by reminders. Therapy widens the window and teaches the return journey.

Points worth keeping

- CPTSD in ICD-11
- Affect dysregulation and self-concept
- Relational disturbance
- Why prolonged trauma differs

PRACTICE TASK

Contrast PTSD and CPTSD in one paragraph.

CHAPTER 2

The phased approach

Complex trauma, encompassing prolonged, repeated, or interpersonal victimisation often commencing in early childhood, demands a treatment architecture that recognises the profound alterations in neurobiological, affective, and relational systems. The most influential articulation of such an architecture is found in Judith Herman's seminal work Trauma and Recovery (1992), which proposes a three-phase model: safety and stabilisation, processing, and integration and reconnection. Subsequent clinicians and researchers, among them Bessel van der Kolk (2014) and Marylene Cloitre, whose survey of expert clinicians found broad support for phase-based treatment of complex PTSD (Cloitre et al., 2011), have refined this phased approach and applied it to Complex PTSD (C-PTSD) and other specialist trauma populations such as survivors of human trafficking, refugees, and individuals with developmental trauma disorder, although the evidence that a phased approach is necessary remains contested (De Jongh et al., 2016).

The neurobiological rationale for a phased approach rests on the concept of the window of tolerance, described by Siegel (1999), which is often read alongside the polyvagal theory of Porges (1995), although the physiological premises of that theory are contested (Grossman, 2023). Traumatic stress can constrict the window, rendering the client either hyper-aroused (fight-flight) or hypo-aroused (freeze-dissociation). When the nervous system remains outside a tolerable range, exposure to trauma-related cues can precipitate a dysregulation spiral, leading to heightened cortisol, fragmented memory encoding, and impaired executive functioning. Consequently, the therapist must first facilitate a state in which the client can self-regulate, maintain affective stability, and engage in reflective cognition before any direct confrontation with traumatic memories is attempted.

Phase one, safety and stabilisation, is therefore the cornerstone of the entire therapeutic trajectory. Safety is multifaceted, encompassing physical safety, emotional safety within the therapeutic relationship, and relational safety in the client's broader environment. The therapist cultivates a secure base through consistent empathy, transparent boundaries, and collaborative

goal-setting, thereby counteracting the attachment ruptures that often accompany complex trauma. A pivotal component of this phase is the development of social support: the systematic identification, validation, and utilisation of the client's existing support networks while concurrently teaching discrimination between safe and unsafe interpersonal contexts. This work often involves mapping the client's relational landscape, rehearsing assertive communication, and establishing contingency plans for crisis situations.

Affect regulation skills are introduced through a repertoire of grounding, mindfulness, and somatic techniques. Grounding exercises, such as the 5-4-3-2-1 sensory method, help the client anchor attention to the present moment, thereby expanding the window of tolerance. Mindfulness practices, adapted from Kabat-Zinn (1994) and tailored to trauma-sensitive protocols, foster non-judgmental awareness of internal states without triggering re-experiencing. Somatic interventions, including diaphragmatic breathing, progressive muscle relaxation, and sensorimotor psychotherapy (Ogden, Minton and Pain, 2006), target the autonomic nervous system directly, promoting parasympathetic activation.

Readiness to transition from phase one to phase two is not a function of elapsed time alone but a qualitative shift in the client's internal capacity. The therapist conducts ongoing assessment using both subjective and objective markers. Standardised measures such as the PHQ-9 and GAD-7 provide quantifiable indices of depressive and anxiety symptomatology, while the International Trauma Questionnaire (ITQ) offers a specific gauge of C-PTSD symptom clusters. A reduction in PHQ-9 and GAD-7 scores, coupled with a stable or decreasing ITQ score for the affective dysregulation items, suggests that the client can tolerate brief exposures to affect-laden material without destabilising. Additionally, the therapist observes behavioural indicators: the client's ability to request a break, employ self-soothing strategies autonomously, and maintain a coherent narrative during session reflections.

Phase two, processing, shifts the therapeutic focus to the traumatic memory itself. The central aim is to transform fragmented, sensory-laden recollections into an integrated autobiographical narrative that no longer dominates present functioning. This phase draws upon evidence-based modalities such as prolonged exposure (Foa et al., 2007), narrative exposure

therapy (Neuner et al., 2004), and cognitive processing therapy (Resick et al., 2017). The therapist selects techniques that align with the client's developmental history, cultural context, and current neurocognitive capacity. For instance, a client with significant dissociative symptoms may benefit from a paced exposure protocol that interleaves imaginal exposure with grounding anchors, whereas a client with strong cognitive distortions may require intensive cognitive restructuring before any imaginal work.

During processing, the therapist facilitates the identification and modification of maladaptive beliefs that have arisen from the trauma, often termed "stuck points" in cognitive models. These beliefs, such as "I am unlovable" or "The world is unsafe," are examined through Socratic questioning, behavioural experiments, and schema-focused interventions. Simultaneously, exposure techniques allow the client to revisit the memory in a controlled manner, diminishing the conditioned fear response via extinction learning. The therapist monitors physiological markers—heart rate variability, skin conductance—and subjective distress ratings (e.g., Subjective Units of Distress Scale) to ensure the client remains within the expanded window of tolerance.

The transition to phase three, integration and reconnection, is predicated on the client's demonstrated ability to recall the trauma without overwhelming dysregulation, to articulate a coherent narrative, and to exhibit adaptive coping in the face of residual triggers. Integration involves consolidating the newly processed memory into the client's broader life story, thereby restoring a sense of continuity and agency. Reconnection focuses on rebuilding roles, relationships, and aspirations that may have been eclipsed by the trauma. This phase aligns with the recovery-oriented principles advocated by the British Association for Counselling and Psychotherapy (BACP) and the NHS's recovery model, emphasising hope, empowerment, and community participation.

In practice, the therapist supports the client in redefining identity beyond the victim role, fostering self-compassion, and engaging in meaningful activities that reinforce a sense of mastery. Techniques such as strengths-based formulation, vocational counselling, and relational repair work are employed. The therapist also revisits social support, encouraging the client to deepen existing supportive bonds and to seek new connections that reflect the client's evolving self-concept.

Judging readiness to move between phases remains a dynamic, iterative process. The therapist must balance the client's expressed desire for progress with objective evidence of stability. If, during phase two, the client experiences a resurgence of panic, severe dissociation, or a marked increase in PHQ-9 scores, the therapist must promptly revert to phase one interventions, reinforcing grounding and affect regulation before resuming processing. Conversely, premature advancement to phase three without sufficient consolidation can result in relapse, as unresolved trauma fragments may re-emerge under stress. The therapist therefore maintains a flexible stance, ready to oscillate between phases as the client's neurobiological and psychosocial landscape evolves.

To illustrate the nuanced application of the phased approach, consider the case of Emily, a 34-year-old woman referred for Complex PTSD following chronic childhood emotional and physical abuse, compounded by adolescent exposure to domestic violence. Emily presented with pervasive hypervigilance, frequent dissociative episodes, a PHQ-9 score of 18, a GAD-7 score of 15, and an ITQ profile indicating severe affective dysregulation and negative self-concept. In the initial phase, the therapist prioritised establishing safety. Through collaborative safety planning, Emily identified a trusted sister and a community support group as primary sources of social support. Grounding techniques, including the use of a tactile stone and paced breathing, were introduced and rehearsed daily. Over a period of ten weeks, Emily's PHQ-9 reduced to 12 and GAD-7 to 9, while her dissociative episodes decreased from daily to twice weekly. Importantly, Emily began to voluntarily employ grounding strategies when she sensed rising anxiety, signalling a shift in internal regulation.


Having demonstrated a stable window of tolerance, the therapist assessed readiness for phase two. Emily reported a willingness to explore the memories of her childhood abuse, yet expressed fear of being overwhelmed. The therapist therefore designed a paced imaginal exposure schedule, beginning with a brief, 5-minute narrative of a neutral childhood event, followed by a grounding anchor. As Emily's distress ratings remained below 4 on the Subjective Units of Distress Scale, the therapist incrementally extended the exposure to include the first instance of emotional abuse. Concurrently, cognitive restructuring targeted Emily's entrenched belief "I am worthless," challenging it with evidence of her academic achievements and supportive relationships. Across twelve weeks of processing, Emily's ITQ

scores for re-experiencing and negative beliefs decreased markedly, and her capacity to discuss the trauma without dissociation improved.

Transitioning to phase three, the therapist facilitated integration by helping Emily write a chronological narrative that linked early abuse experiences with later relational patterns, thereby creating a cohesive autobiographical account. Reconnection work involved exploring Emily's long-standing interest in horticulture. With the therapist's support, Emily enrolled in a community gardening project, re-established contact with a former colleague, and set a goal to volunteer once weekly. Over the subsequent eight weeks, Emily's PHQ-9 fell to 6, GAD-7 to 4, and she reported a renewed sense of purpose and belonging. Her social support expanded as she cultivated new friendships within the gardening group, illustrating the reciprocal reinforcement between integration and reconnection.

Emily's trajectory exemplifies the essential principles of the phased approach: the necessity of establishing safety and affect regulation before confronting trauma, the careful calibration of exposure based on physiological and subjective markers, and the ultimate goal of restoring a coherent identity that can engage meaningfully with the world. The therapist's ongoing assessment of readiness, guided by validated measures and real-time observation, ensured that each transition was anchored in genuine capacity rather than arbitrary timelines.

In sum, the phased approach to Complex PTSD and specialist trauma populations remains a robust, evidence-informed framework that aligns neurobiological imperatives with therapeutic pragmatism. Phase one lays the groundwork of safety, affect regulation, and social support; phase two facilitates the controlled processing of traumatic memories through exposure and cognitive restructuring; and phase three consolidates these gains into an integrated self-concept and renewed community participation. Judging readiness to move between phases demands a vigilant, data-driven stance that respects the client's lived experience while safeguarding against destabilisation. As the field continues to evolve, emerging research on neurofeedback, sensorimotor psychotherapy, and culturally adapted interventions will likely enrich each phase, yet the core principle—that the client must first be able to stay within a tolerable window of arousal before the trauma can be safely revisited—will endure as the keystone of effective trauma-focused care.



Phases 3–6 repeat per target memory; the client's own processing does the work.

FIGURE 2.1 The eight phases of EMDR

Shapiro's standard protocol moves through eight phases. Preparation and history-taking do the stabilising groundwork; assessment activates the target memory with its image, cognitions and body sensation; desensitisation applies bilateral stimulation until distress falls; installation and body scan consolidate; closure and re-evaluation keep every session safe and cumulative.

Points worth keeping

- Phase one: safety and stabilisation
- Phase two: processing
- Phase three: integration and reconnection
- Judging readiness to move between phases

PRACTICE TASK

Outline what phase-one work looks like for a complex presentation.