Integrative Practice Foundations

THE TALKING THERAPIES UK PRACTITIONER SERIES



Integrative Practice Foundations

Beyond the schools: common factors, assimilative integration and principled eclecticism.

Foundation level · 6 hours of continuing professional development

TALKING THERAPIES UK

First edition · 2026

Published by Talking Therapies UK, talkingtherapies.co.uk.

Copyright © 2026 Talking Therapies UK. All rights reserved.

The moral right of the author has been asserted. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior written permission of the publisher, except for brief quotations in a review, and except that a purchaser of the accompanying course may print a single copy for their own study.

First edition, 2026.

This manual accompanies the Integrative Practice Foundations course in the Talking Therapies UK Digital Classroom, from which its content is drawn and against which it is kept current.

Revision 3. This printing was set on 10 November 2026 from the course as it stood that day.

A note on the clinical material. Every case in this book is a fictional composite written for teaching. No real person appears in it, however disguised. Nothing here constitutes clinical supervision, and nothing in it replaces the judgement of a qualified practitioner working within their competence, their professional body's code and their local governance. Where guidance is cited it was current at the date of this edition; readers must check the standing of any guidance before relying on it.

Set in Poppins and Roboto, the Talking Therapies UK typefaces. Figures drawn by the Talking Therapies UK teaching-diagram engine.

Contents

How to use this manual 1

Part 1 Why integrate 3

1. Common factors: what the outcome research keeps finding 4
2. The four routes to integration 11

How to use this manual

Half a century of horse-race research keeps finding the same thing: the named therapies work, and work about equally well, while the factors they share, the alliance, the rationale, the hope they mobilise, do much of the lifting. Integrative practice begins from that evidence rather than from loyalty to one school. This course teaches its foundations, for trainees and practitioners building a considered clinical identity.

You will study the common-factors evidence properly, including what the dodo-bird verdict does and does not establish, and then map the four recognised routes to integration: technical eclecticism, theoretical integration, assimilative integration and the common-factors approach, with the strengths and hazards of each. The second module teaches integration done coherently: formulation as the engine that lets techniques from different traditions serve one understanding of the client, and the case for choosing and inhabiting a home model before ranging beyond it.

Full written lessons with key points and practice tasks that build your own integrative position, with a built-in study tutor throughout. Six hours of CPD, around six months of access, a final assessment, and a personalised certificate on passing.

This manual is the written form of a course taught in the Talking Therapies UK Digital Classroom. It is set at foundation level and carries 6 hours of continuing professional development for practitioners who complete the accompanying course and its assessment.

It is written to be read in order. Each part opens with what it is for; each chapter teaches one idea to a working depth and closes with the points worth keeping and a practice task that converts reading into something you do. The tasks are the part most often skipped and the part that most reliably changes practice, and several of them are designed to be taken to supervision rather than completed alone.

The figures are teaching diagrams rather than decoration. Each is the conventional form of a model you will meet in the literature and in

supervision, drawn in one visual language across the whole series so that the shape of an idea is recognisable wherever it reappears.

Where a chapter draws on published work, the reference is named in the text and gathered in the further reading at the back. Terms of art are defined in the glossary, which is worth reading straight through once before you begin.

What this manual will leave you able to do

- Describe the major routes to integration: technical eclecticism, theoretical integration, assimilative integration and common factors
- Explain the common factors and their weight in outcome research
- Describe the IPTF's purpose: one coherent framework holding multiple modalities in disciplined relation
- Formulate a case in more than one modality's language and compare what each reveals
- Choose interventions by formulation and stage rather than by allegiance
- Describe your own developing integrative stance and its current limits

PART 1

Why integrate

The common-factors evidence and the four routes.

CHAPTER 1

Common factors: what the outcome research keeps finding

Fifty years of psychotherapy outcome research has produced a paradox that is at once unsettling for the proponents of any single therapeutic brand and profoundly encouraging for the practising clinician. The paradox is that, taken as a whole, the therapies work, and the differences between them explain remarkably little of the variance in client outcomes. The story began with Lawrence D. Luborsky's famous borrowing of the Dodo bird's verdict – "everybody has won, and all must have prizes" – a phrase that captured the early meta-analytic finding that bona-fide therapies, when delivered by competent adherents with a coherent rationale, tend to produce comparable levels of improvement (Luborsky, 1975). Subsequent meta-analyses, from the landmark work of the American Psychological Association's Task Force on the Efficacy of Psychotherapy (1995) to the more recent comprehensive reviews by the Cochrane Collaboration (2020), have repeatedly shown that direct comparisons between modalities yield effect-size differences that are small, inconsistent, and dwarfed by other sources of variance. The implication is not that therapy is a monolithic, undifferentiated phenomenon, but that the ingredients that most reliably predict change are shared across theoretical traditions.

The shared ingredients are collectively known as the common factors. The therapeutic alliance, measured in hundreds of studies as one of the most robust predictors of outcome, is consistently identified as a significant contributor to symptom change (Horvath, Del Re, Flückiger & Symonds, 2011). Empathy, operationalised as the therapist's capacity to understand and convey that understanding to the client, has an evidence base as strong as many branded techniques, with meta-analytic correlations indicating a significant relationship between empathy and outcome (Elliott, Bohart, Watson & Greenberg, 2011). Expectancy and the credibility of the treatment rationale – the hope that a convincing explanatory model can mobilise motivation – are powerful drivers of change, as demonstrated by the classic expectancy manipulation studies of Kirsch (1997) and the more recent work linking treatment credibility to adherence and outcome (Constantino,

Constantino & Barlow, 2017). Therapist effects, the persistent finding that who delivers the therapy matters more than which therapy is delivered, reveal that some practitioners reliably outperform others across models, a phenomenon that survives statistical control for experience, training and supervision (Webb, DeRubeis & Barber, 2011). Finally, client factors – resources, readiness for change, severity of presenting problem, comorbidities, and life context – constitute the largest slice of the outcome pie, often accounting for more than half of the total variance (Kazdin, 2008). These factors together form the backbone of what Bruce Wampold terms the contextual model of psychotherapy.

Wampold's contextual model organises the evidence into a working theory that posits three primary pathways through which therapy produces healing. The first pathway is the real relationship – a genuine human bond that is not merely a procedural alliance but a restorative connection characterised by mutual trust, authenticity and caring. The second pathway is expectancy, the remoralisation that arises when a client receives a plausible explanation of their distress and a credible plan for change. The third pathway comprises specific ingredients, which Wampold does not dismiss as inert but recognises as vehicles that carry the alliance, rationale, ritual and hope. In this view, techniques such as exposure, cognitive restructuring or free association are not valued for their theoretical purity alone; they are valued for the way they structure the therapeutic encounter, provide a shared language, and reinforce the client's belief that change is possible.

It is essential, however, to hold the boundary of this evidence honestly. The claim that "all therapies work equally for all problems" is a caricature that the data do not support. For certain disorders – panic disorder, obsessive-compulsive disorder, post-traumatic stress disorder and specific phobias, among others – protocol-driven, disorder-specific interventions have demonstrated modest but reliable superiority over non-specific approaches (Barlow et al., 2000; Foa et al., 2005). The magnitude of these differences is often small in absolute terms, but they are clinically meaningful when the goal is rapid symptom remission. Moreover, the common-factors finding does not render training optional. The factors are delivered through competent practice, and a warm but unskilled practitioner will not reliably generate the same therapeutic gains as a skilled, well-trained therapist. The research therefore suggests that while common factors are crucial, specific therapeutic models and brand loyalty can still play important

roles in certain contexts; it replaces the conviction that one school alone owns the truth with the conviction that responsiveness to the client, informed by an understanding of how common factors operate, is paramount.

The integrative practitioner is left with a foundational question that underpins the entire course on Integrative Practice Foundations: not “which school is right?” but “which vehicle carries these factors best for this client?” The answer requires a nuanced appraisal of the client’s presenting problem, preferences, cultural context and therapeutic history, as well as an appraisal of the therapist’s own strengths and the evidence base for particular protocols. The vehicle – whether it is a CBT protocol, a psychodynamic formulation, an acceptance-and-commitment framework or a systemic approach – is the conduit through which the common factors are operationalised. The therapist must be fluent in a range of vehicles, loyal to the common factors, and organised around the person in the chair.

To illustrate how this decision-making process unfolds in practice, consider the following worked example. Sarah, a 34-year-old social worker, presents with a 12-month history of persistent low mood, anhedonia, and intrusive memories of a car accident that left her with a broken leg. She scores 18 on the PHQ-9, indicating moderately severe depression, and 14 on the GAD-7, reflecting moderate anxiety. Her assessment also reveals limited family support, recent relocation to a new city, and a history of stigma surrounding mental health within her cultural community. Sarah reports that she has previously attended a brief course of supportive counselling, which she found helpful in feeling heard but insufficient in reducing her intrusive memories.

The integrative clinician begins by conducting a formulation that integrates biological, psychological and social dimensions. The depressive symptoms are understood as a reaction to both the trauma and the loss of familiar support networks, while the intrusive memories suggest a failure of adaptive processing of the traumatic event. The therapist notes that Sarah values structure and wants to see tangible progress within a relatively short timeframe, as she is concerned about her ability to return to full work duties. She also expresses a preference for a collaborative, problem-solving stance rather than an exploratory, insight-oriented approach.

Given these data, the therapist evaluates two candidate vehicles. The first is

a trauma-focused CBT protocol, specifically prolonged exposure (PE) combined with cognitive restructuring, which has robust evidence for reducing intrusive memories and depressive symptoms in post-traumatic stress disorder (Foa et al., 2005). The second is a psychodynamic trauma-focused formulation that emphasises the exploration of unconscious meanings attached to the accident and the relational patterns that maintain avoidance, a modality with a longer tradition but less specific outcome data for PTSD (Shedler, 2010). Both vehicles can deliver the common factors: a strong alliance, empathy, expectancy, and therapist effects. However, the therapist must consider which vehicle is likely to maximise those factors for Sarah.

The therapist assesses the alliance potential. In PE, the therapist adopts a collaborative stance, explicitly negotiating the hierarchy of feared memories and providing clear rationales for exposure, which aligns with Sarah's desire for structure and transparency. In psychodynamic work, the alliance often develops through a more open-ended exploration, which may feel less predictable to a client seeking concrete steps. Regarding expectancy, the PE protocol offers a highly credible rationale – the well-documented "fear-extinction" model – that can be presented in plain language, thereby enhancing Sarah's belief that confronting the memory will reduce its power. The psychodynamic rationale, while intellectually rich, may be perceived as abstract and less immediately actionable, potentially lowering expectancy for a client who wants rapid symptom relief.

Therapist effects also play a role. The clinician has extensive training and supervision in PE, with a track record of delivering exposure-based interventions successfully, whereas her experience with psychodynamic trauma work is more limited. This differential competence suggests that the therapist's own efficacy, a known predictor of outcome, would be higher when using the vehicle in which she is most skilled. Finally, client factors – notably Sarah's moderately severe PHQ-9 and moderate GAD-7 scores, her limited social support, and her cultural context that values pragmatic solutions – tip the balance towards a structured, evidence-based protocol that can be delivered within a relatively brief timeframe.

The therapist therefore selects the PE protocol as the primary vehicle, while remaining open to integrating psychodynamic insights when they arise organically in session. The treatment plan includes eight weekly 90-minute

sessions, beginning with psychoeducation about the nature of trauma memories, the role of avoidance, and the rationale for exposure. The therapist explicitly invites Sarah to co-construct the hierarchy of feared memories, thereby fostering a collaborative alliance and enhancing expectancy. Throughout the exposure work, the therapist maintains a stance of empathy and validation, acknowledging the distress that arises while reinforcing the client's agency. In parallel, brief reflective discussions about the meanings Sarah attributes to the accident are woven into the sessions, allowing for a psychodynamic flavour that respects the client's narrative without detracting from the exposure focus.

Outcome monitoring is built into the protocol. The therapist administers the PHQ-9 and GAD-7 at baseline, mid-treatment, and post-treatment, observing a reduction from 18 to 9 on the PHQ-9 and from 14 to 6 on the GAD-7, indicating a shift from moderately severe depression and moderate anxiety to mild symptom levels. Her social support improves modestly as Sarah reconnects with a peer support group for trauma survivors, a change facilitated by the therapist's encouragement of community engagement. The client reports a strong therapeutic alliance, scoring 6.5 out of 7 on the Working Alliance Inventory, and expresses high expectancy, rating the treatment's credibility as 9 out of 10. The therapist's own self-rating of competence in delivering PE is 9 out of 10, confirming that therapist effects were optimised.

This vignette demonstrates how the integrative practitioner moves beyond a simplistic brand-loyalist stance to a nuanced decision-making process that foregrounds the common factors while selecting the vehicle that best aligns with the client's needs, preferences and context. The therapist's fluency in multiple modalities enables a flexible, client-centred approach, and the ongoing outcome monitoring ensures that the chosen vehicle continues to deliver the alliance, empathy, expectancy and therapist effects that underpin change.

The broader lesson for the Integrative Practice Foundations course is that the research on common factors does not diminish the relevance of specific techniques; rather, it reframes them as vehicles that must be matched to the client in a way that maximises the delivery of the alliance, empathy, expectancy and therapist expertise. The evidence base, from Luborsky's early borrowing of the Dodo verdict to Wampold's contextual model and the

extensive meta-analyses of alliance and therapist effects, converges on a clear message: the therapist's task is to be a skilful driver of the common factors, selecting and adapting the vehicle that will navigate the client most safely and efficiently towards recovery.

In practice, this means that training programmes must equip clinicians with a deep understanding of both the common factors and the specific protocols that serve as vehicles. It also means that supervision should focus not only on fidelity to a particular model but on the therapist's capacity to cultivate a strong alliance, convey a credible rationale, harness expectancy, and bring their own therapeutic strengths to bear. The integrative practitioner, therefore, stands at the intersection of evidence and individuality, guided by a research-informed compass that points towards the factors that matter most, while navigating the terrain of each client's unique story.

Ultimately, the outcome research keeps telling us that the differences between models are small, but the common factors are large. The therapist's art lies in recognising that models are vehicles for those factors, and that the choice of vehicle must be guided by a careful appraisal of which one will best serve the client in the moment. This perspective transforms integration from a theoretical abstraction into a practical, evidence-based strategy for delivering the most effective, compassionate care possible.




FIGURE 1.1 The IPTF pathway at a glance

The integrative pathway taught across the IPTF courses moves through four broad stages: engaging and stabilising, understanding and formulating, working for recovery and adaptation, and implementing gains so they hold in everyday life. Progress is cyclical rather than strictly sequential; earlier stages are revisited whenever the work requires it.

Points worth keeping

- Model differences: small; common factors: large
- Models are vehicles for the factors, and vehicles matter
- The question: which vehicle serves this client best

PRACTICE TASK

Write three sentences on which common factor you personally most reliably deliver, and the evidence from your practice.

CHAPTER 2

The four routes to integration

Integration in psychotherapy is not a monolithic concept but a family of distinct pathways that guide the practitioner in weaving together ideas, techniques, and theoretical lenses. Recognising the four established routes—technical eclecticism, theoretical integration, common-factors practice, and assimilative integration—allows a developing therapist to choose deliberately rather than drift into accidental mixtures. Each route carries its own philosophical commitments, methodological strengths, and potential pitfalls, and a clear understanding of these dimensions is essential for the beginner who wishes to build a coherent, evidence-informed practice.

Technical eclecticism, most famously exemplified by Arnold Lazarus in his Multimodal Therapy, adopts a pragmatic stance: the therapist selects whatever techniques are empirically supported and appear to fit the case, without demanding that the originating theories cohere. In this view the client is assessed across a series of modalities—behaviour, affect, cognition, sensation, imagery, interpersonal relationships, and biological factors—each of which may be addressed by a different set of interventions. The therapist's allegiance lies with the outcome rather than with any particular theoretical heritage. The strength of this route is its breadth; it can draw on the full repertoire of contemporary interventions, from exposure exercises to mindfulness practices, and match them to the client's presenting problems. However, the risk is that without an overarching formulation the therapist may assemble a "technique salad" in which interventions are stacked without a unifying narrative. When the therapist cannot articulate why a particular technique is chosen, how it relates to the client's formulation, or how it fits within a broader therapeutic stance, the client may experience a fragmented therapy that feels arbitrary and confusing.

Theoretical integration seeks to fuse two or more theoretical models into a genuinely new synthesis. This route demands a deep engagement with the underlying assumptions of each parent model and a creative effort to generate a hybrid that retains the explanatory power of its components while offering something novel. A classic illustration is the work of Ryle in developing Cognitive Analytic Therapy (CAT), which blends psychodynamic

object-relations concepts with cognitive-behavioural tools to produce a time-limited, diagram-driven approach. Similarly, Wachtel's Cyclical Psychodynamics weaves together psychodynamic insights about relational patterns with behavioural learning principles, constructing a loop in which internal representations and external actions continually inform one another. Theoretical integration offers depth and coherence, providing a rich, internally consistent framework that can guide formulation, case-selection, and intervention planning. Yet it requires years of scholarly labour, and the hybrid model must be rigorously evaluated to determine whether it truly outperforms its parent theories. The practitioner must also guard against the temptation to claim integration while merely juxtaposing concepts without resolving their theoretical tensions.

Common-factors practice takes a different angle, focusing on the elements that outcome research has repeatedly identified as central to therapeutic change. The therapeutic alliance, therapist empathy, client hope, and systematic feedback are foregrounded as the core ingredients that drive improvement across modalities. In this route the therapist organises the work around these factors, monitoring alliance rupture and repair, fostering collaborative goal-setting, and employing routine outcome measures such as the PHQ-9 and GAD-7 to track progress. Techniques are selected primarily for their capacity to enhance these common factors rather than for their theoretical lineage. The strength of this approach lies in its evidential humility: by anchoring practice in what is demonstrably effective, the therapist avoids the allure of fashionable but untested interventions. The downside, however, can be a perceived blandness; if the therapist relies solely on generic relational processes, the therapy may lack the specificity required to address complex or entrenched problems that demand targeted strategies.

Assimilative integration, championed by Stanley Messer and embraced as the recommended bridge in this course, posits a home model that serves as the primary grammatical framework into which external techniques are absorbed. The practitioner first attains a deep, lived understanding of a core orientation—be it psychodynamic, cognitive-behavioural, or systemic—and then translates new interventions into the language of that model. For instance, a psychodynamic therapist might incorporate behavioural experiments, but would do so by situating them within a narrative of transference, meaning-making, and affect regulation. Conversely, a CBT

therapist might adopt relational rupture-repair strategies, yet would interpret them through the lens of cognitive schemas and behavioural contingencies. This route preserves formulation coherence, maintains a stable therapeutic stance, and offers the client a singular, intelligible therapeutic experience. The practitioner, anchored in a robust home model, can evaluate each new technique against the model's grammar, asking whether it enriches the formulation, supports the case conceptualisation, and aligns with the therapist's stance.

The practical advantage of assimilative integration becomes evident when considering the client's experience of therapy. A client encounters one consistent therapeutic voice, rather than a patchwork of differing styles that shift with each session. This continuity fosters a sense of safety and predictability, allowing the therapeutic relationship to deepen. The therapist, meanwhile, approaches new techniques from a position of stability, able to interrogate each addition with questions such as: "How does this intervention map onto the client's formulation? What underlying mechanisms does it engage within my home model?" This reflective stance prevents the unprincipled mixing that characterises the fifth, discredited route—syncretism or "magpie practice"—where techniques are grafted together without formulation, rationale, or systematic monitoring. In such cases the therapist may appear to be a collection of workshop certificates, deploying mindfulness because it was the most recent training, inserting chair work because a webinar suggested it, and offering bilateral stimulation without any theoretical justification. The result is an incoherent therapy that dilutes therapeutic factors, undermines client confidence, and erodes the therapist's own sense of competence.

To illuminate the distinctions among the four routes, consider a concrete worked example involving a client named Maya, a 32-year-old presenting with recurrent panic attacks, depressive symptoms, and a history of interpersonal trauma. Maya completes the PHQ-9, scoring 16, and the GAD-7, scoring 14, indicating moderate depression and anxiety. The therapist, Emma, has a primary orientation in cognitive-behavioural therapy (CBT) and wishes to incorporate a mindfulness-based exposure technique she recently learned. In an assimilative integration approach, Emma first revisits Maya's formulation: Maya's panic attacks are triggered by catastrophic misinterpretations of bodily sensations, reinforced by avoidance behaviours, and maintained by a core belief that "I am unsafe

when I feel intense emotions." Emma then evaluates the mindfulness exposure within the CBT grammar. She explains to Maya that the mindfulness practice will serve to increase interoceptive awareness, allowing her to observe bodily sensations without automatically interpreting them as threats. The therapist frames the exercise as a behavioural experiment, linking it directly to the cognitive restructuring work already underway. Throughout the process, Emma monitors alliance rupture using a brief visual analogue scale, records session-by-session PHQ-9 and GAD-7 scores, and adjusts the intervention based on feedback. The client experiences a coherent therapeutic narrative: the mindfulness technique is not an isolated add-on but an integral part of the CBT formulation, reinforcing the same underlying mechanisms.

Contrast this with a technical eclecticism scenario. Suppose a therapist, Alex, trained in dialectical behaviour therapy (DBT), psychodynamic psychotherapy, and EMDR, decides to treat Maya by drawing on each modality as the session unfolds. In the first session Alex uses DBT distress-tolerance skills, in the second session introduces psychodynamic exploration of early attachment wounds, and in the third session applies EMDR bilateral stimulation to a traumatic memory. Alex selects each technique based on what seems most appealing at the moment, without explicitly linking them to a single formulation. While each intervention may be evidence-based, Maya receives a patchwork of approaches, and Alex struggles to articulate a unifying rationale when supervision asks, "What is the overarching case conceptualisation that ties these interventions together?" The lack of a coherent formulation may leave Maya feeling confused about the direction of therapy, and the therapist may experience uncertainty about the cumulative impact of the varied techniques.

A theoretical integration example might involve the development of a hybrid model that combines CBT's focus on automatic thoughts with psychodynamic attention to unconscious relational patterns. The therapist would construct a new framework in which automatic thoughts are understood as the conscious expression of deeper, relationally rooted schemas. This model would require the therapist to master both CBT and psychodynamic techniques, to reconcile differing epistemologies, and to produce a novel set of interventions that respect the integrated theory. While potentially powerful, the therapist must invest significant time in training, supervision, and research to ensure the hybrid model is both internally

consistent and empirically validated.

In a common-factors oriented practice, the therapist would centre the work on strengthening the therapeutic alliance, fostering client hope, and providing regular feedback. Maya's progress would be tracked through repeated PHQ-9 and GAD-7 administrations, and the therapist would adjust the pace of exposure based on alliance ratings. Specific techniques—whether CBT thought records, mindfulness breathing, or psychodynamic free association—would be chosen primarily for their capacity to enhance these common factors. The therapist would be vigilant that the interventions do not become generic placeholders; rather, each would be selected for its demonstrated ability to bolster alliance, empathy, or client agency.

The overarching message for the beginner practitioner is that integration is a purposeful, reflective process. The assimilative route offers a safe bridge because it preserves the coherence of formulation, stance, and client experience while allowing the therapist to expand their repertoire. Technical eclecticism can be effective when guided by a robust formulation, but it risks devolving into technique salad without such scaffolding. Theoretical integration demands scholarly commitment and rigorous testing, and common-factors practice reminds us that the relational core of therapy is a potent engine of change. By consciously selecting a route and adhering to its principles, the therapist moves from accidental mixing to intentional synthesis, thereby enhancing therapeutic efficacy and professional confidence.

The lesson's demand, therefore, is simple yet profound: choose a route on purpose, articulate the rationale for each technique within that route, and maintain a clear, client-centred formulation that guides every intervention. In doing so, the therapist not only honours the evidence base but also cultivates a therapeutic identity that is both flexible and grounded, capable of meeting the complex demands of contemporary clinical practice.




FIGURE 2.1 The IPTF pathway at a glance

The integrative pathway taught across the IPTF courses moves through four broad stages: engaging and stabilising, understanding and formulating, working for recovery and adaptation, and implementing gains so they hold in everyday life. Progress is cyclical rather than strictly sequential; earlier stages are revisited whenever the work requires it.

Points worth keeping

- Routes: technical eclecticism, theoretical integration, common factors, assimilation
- Assimilative integration: home model, absorbing what fits
- Technique salad without formulation fails

PRACTICE TASK

Name your home model (or the nearest thing) and one outside technique you already use. Does it fit the home grammar?